

Biological Riches and Rarities



After dropping supplies at the space station, the shuttle circles Earth once, then twice. Soon the crew will fire the thrusters and return home to California. On the planet below, the Sun rises and sets every 16 minutes. Where it is day, the crew notices areas of brilliant green. The oceans are the deepest blue imaginable. Rivers appear as lines, like veins on the back of a grandmother's hand. From here, Earth seems so fragile, a colorful, spinning globe.

California's Biodiversity

Members of the crew cannot help but think about how rich Earth is. It is home to so many different forms of life. Every living thing depends on another, playing an integral part in its own natural community. Many biological communities exist on Earth. California, for example, has 5,879 species of plants and animals (more than any other state in the United States) and 2,214 endemic species—plants and animals found nowhere naturally beyond its borders.

Biodiversity (biological diversity) is a term used to describe the richness in numbers and variety of species of plants and animals in an area or region.

Climate and Geography

California's Mediterranean climate, with hot, dry summers and mild, wet winters, makes it an ideal place for many species to thrive. Because of its rich biological diversity and the threats to many of its species, biologists have designated California a biodiversity "hotspot."



Earth viewed from space

Biodiversity hotspots are home to a rich diversity of Earth's plant and animal species. A combination of factors makes them like "magnets" for life. The nation's largest breeding ground for birds, for example, is in California. The state is also home to 20 species of freshwater fish, 17 reptiles, and 17 mammals that are found in no other place on the planet.

California's biodiversity can also be explained by its unique geography. The state has both the highest point in the continental United States at the summit of Mount Whitney



Mount Whitney, California



Joshua trees, Mojave Desert, California

(14,494 feet) and the lowest point at Badwater, Death Valley (282 feet below sea level). These distinct elevations are only 80 miles apart, and both are less than 200 miles from the Pacific Ocean. The differences in elevation, combined with California's moderate climate and its 100 million acres, create a mosaic of habitats where life has expanded to take advantage of a multitude of opportunities.

Home to Deserts, Forests, Wetlands, and More

As the shuttle nears Earth, the crew can see the wedge shape of the Mojave Desert boldly outlined by the mountains and fault lines that define it. It is considered “high” desert, with an average elevation of 3,500 feet, though it also extends into the lowest corners of Death Valley.

The high desert climate is dry. There is a summer thunderstorm season between July and September, but most rain falls in the winter, and snow is common at higher elevations. The range of temperatures makes it one of the state's greatest centers for biodiversity. Approximately 1,500 plant species live in the

Mojave Desert, with 210 of them identified as endemic.

The scientists in the crew especially love the woody Joshua trees (*Yucca brevifolia*), with “hairy” limbs that seem to stretch for something just beyond their reach. They are endemic in this area. Joshua trees extend roots to take advantage of both surface and deeper sources of water. They are well adapted to frost in the winter and high desert temperatures in the summer.

A number of organisms depend on the Joshua tree for survival, just as the Joshua tree depends on them. One of these is the Tegeticula moth. This small, white moth lays her eggs inside the white blossoms of the Joshua tree, pollinating the flowers, so the embryonic seeds inside will grow. When the larvae of the moth hatch, they feed on a few small seeds. The symbiotic



Carrizo Plain grasslands, California

relationship between the moth and the Joshua tree allows both organisms to produce new life.

As the shuttle drops further, the crew can see beyond the desert to the grasslands of the Carrizo Plain. With the shuttle still far above Earth, the homesick crew talks about how the constant wind plays with sunlight on the natural grasses, tossing them back and forth. Though much of the original grassland of the Great Central Valley has been replaced by agriculture, the protected Carrizo Plain remains largely untouched. It is home to some of California's endangered animals, including the San Joaquin Valley kit fox and the blunt-nosed leopard lizard.

Coral Hollow, in the northwestern San Joaquin Valley, provides another

example of original grassland habitat. A variety of reptiles and amphibians, including the California red-legged frog and the western pond turtle, live there. Composed of 70% grassland and 30% riparian woodland, Coral Hollow also displays unique vegetation, including the desert olive, iodine bush, and honey mesquite.

As they survey the land, crewmembers imagine the wildflowers that must be blooming wherever there is enough water to germinate their seeds. They talk about the vernal pools, seasonal wetlands that occur in the Central Valley



Kit fox cubs

and other regions. These pools support numerous plants and animals, some threatened with extinction. Formed in the winter months when rain fills clay and hardpan (dense layers of soil that do not easily absorb water) hollows with rainwater, vernal pools provide a temporary home to fairy shrimp, frogs,



Wildflowers by vernal pool

and many aquatic plants. In the spring when the water begins to evaporate, colorful wildflowers spring up, lining the pools with chains of goldfields, creamy Johnny tuck, and lacy meadowfoam.


One crewmember recalls traveling through California to the “land of the giants,” the deep old-growth forest that adds so much to biodiversity. Although old-growth redwood forests are wonderful ecosystems, they are known for their remarkable lack of California’s biodiversity compared with other forests. Except for a 14-mile extension into Oregon, the coast redwood forest belongs only to California. Redwoods, Earth’s tallest trees, sometimes stand 300 feet tall and have trunks that reach 25 feet in diameter at their bases. The average redwood is 600 years old, though many are much older. When they fall, redwoods act as their own “nursery logs,” nurturing rows of new seedlings that eventually grow into mature trees. Because of their incredibly thick bark, old-growth redwoods are very fire resistant.

Under the shade of the giant redwood canopy, several trees provide a tall secondary layer of growth. Tanoak, California bay, Douglas fir, hemlock, and sometimes big leaf maple grow with filtered sunlight. Wild

rhododendron, poison oak, salal, and huckleberries fill in the next colorful layer. The ground level is rich with redwood sorrel and wood ferns. They provide a safe hiding place for many animals, including the banana slug, Pacific giant salamander, and the red-legged frog.

It is almost time to land, and the crew directs its attention to the desert below. Their high altitude tour of California’s biodiversity is nearing an end, and so many places remain to visit! Rogers Dry Lake is now directly in front of the shuttle, its drawn runways clearly marked on hard clay.

As the shuttle touches down, the crew looks to the north, where sunlight illuminates the Sierra Nevada Mountain Range. They take deep breaths, glad to



Coast redwood, California

be home. Earth from a distance, though stunning, is nothing compared to the rich beauty and diversity that surround the astronauts when their feet finally touch the ground.



Sierra Nevada range



Bay Area/Delta Bioregion

This bioregion takes its name from the San Francisco Bay and Delta watersheds. It contains freshwater marshes and salt marshes. Hilly coastal areas support coastal prairie scrub,

mixed hardwood, and valley oak communities. Sonoma, Marin, and San Mateo Counties have redwood forests. Temperatures are moderate, and winters are rainy.

Human Social Systems

More than 6.6 million people live here. Only the South Coast bioregion has more people. Cities include: San Francisco, Santa Rosa, Oakland,

Berkeley, Vallejo, Concord, and San Jose.

Businesses include: tourism, high-tech industries, wine making, and banking. Fishing, shipping, and oil refining occur along the coast. Dairy farming, orchards, and vineyards are common. Water from the Sierras flows through reservoirs and canals. This water supplies two-thirds of California's drinking water. Other uses of the water are irrigating crops and maintaining aquatic habitats.

Natural Systems

Hilly areas along the coast support oak woodlands and grasslands. Coastal salt marshes around San Francisco Bay and freshwater marshes in the Delta provide food and shelter to many birds.



Saltwater marsh, San Francisco Bay

- **Habitats:** Prairie scrub, mixed hardwoods, valley oaks, eucalyptus, manzanita, northern coastal scrub, coastal prairie scrub in the Bay Area; mixed hardwoods, valley oaks, redwoods, salt marshes, freshwater marshes in the Delta.
- **Wetland plants:** Pickleweed, great bulrush, saltbush, cattail.
- **Common birds:** Canvasback, western grebe, black-crowned night heron, great egret, snowy egret, California brown pelican, white pelican, seagull, acorn woodpecker, golden eagle, western bluebird, Caspian tern, American avocet, cedar waxwing.
- **Common mammals:** Grey fox, mule deer, bobcat, raccoon.
- **Other animals:** Pacific tree frog, swallowtail butterfly, painted lady butterfly.
- **Marine life:** Chinook salmon, harbor seal, sea lion, leopard shark, bat ray.
- **Endangered species:** California least tern, California black rail and clapper rail, Smith's blue butterfly, salt marsh harvest mouse, San Francisco garter snake, California freshwater shrimp, northwestern pond turtle, tidewater goby (fish). Rare plants include Marin western flax, Baker's manzanita, Point Reyes checkerbloom, Sonoma sunshine.



Muir Woods, Marin County, California



Clapper rail


Factors Affecting Wildlife Diversity

- habitat loss due to urban, residential, and agricultural development
- water management issues including reduced water for wildlife
- water pollution
- invasive species

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Bobcat



San Francisco area traffic



Central Coast Bioregion

This coastal region features beaches, dunes, and mountains.

Plant communities include chaparral, mixed hardwood and redwood forests, and oak woodlands. The climate is mild

and sometimes foggy. Winters are moist.

Human Social Systems

Artichokes, garlic, and a wide range of fruit and vegetable crops grow in this region. Dairy and cattle ranches are common. Other industries include tourism, wine making, fishing, and oil production. Development pressure is intense, but large blocks of natural land offer many opportunities for conservation.

Natural Systems

Coastal communities bring together marine, freshwater, and terrestrial systems.

- **Mixed coniferous forests in the coastal ranges:** Ponderosa pine, Douglas fir, red alder, and redwoods in the north.
- **Oak woodlands:** Coast live oak, valley oak.
- **Rare tree species:** Monterey pine, Santa Lucia fir.
- **Common birds:** Western snowy plover, willet, whimbrel, long-billed curlew, marbled godwit, American avocet, peregrine falcon, golden eagle, and California spotted owl.
- **Common mammals:** Mountain lion, bobcat, American badger.



Central coast, California

Inland mountain ranges have drier climates and support oak woodlands, grasslands, interior chaparral, and desert-like interior scrub. Oak woodland communities support more than 200 species of plants, 300 vertebrates, and 5,000 invertebrates.

- **Common mammals:** Western gray squirrel, dusky-footed woodrat, Monterey dusky-footed woodrat, pallid bat, Townsend's big-eared bat.

Grasslands now contain mostly nonnative grasses.

- **Rare birds:** Burrowing owl.
- **Common mammals:** California ground squirrel, black-tailed jackrabbit. Tule elk and pronghorn have been reintroduced in the southern part of the region.
- **Rare mammals:** Giant kangaroo rat, San Joaquin kit fox, American badger.

Interior chaparral has a hot, dry climate. It supports drought-resistant shrubs, such as manzanita, California lilac, and chamise.

- **Endangered species:** Morro Bay kangaroo rat, Pacific pocket mouse, California brown pelican, California condor, Santa Cruz long-toed salamander, unarmored threespine stickleback (fish), San Francisco garter snake, southern sea otter, leatherback turtle.

Factors Affecting Wildlife Diversity

- water management conflicts and water transfer
- inappropriate off-road vehicle use
- loss and degradation of dune habitats
- disruption of sand transport processes
- invasive plant and animal species
- growth and development of communities

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Coastal Redwoods, California



Southern sea otter



Pelicans, willets, and godwits



Artichoke farmer



Colorado Desert Bioregion

This bioregion forms part of the larger Sonoran Desert, which reaches across several southwestern states. The Colorado Desert is lower and flatter than the Mojave Desert to the north. It has higher summer daytime temperatures. In winter, frost is rare. Southern sections have rainy seasons in the winter and late summer.

Human Social Systems

Compared with most other bioregions, few people live in the Colorado Desert. Cities include: Palm Springs, Rancho Mirage, El Centro, Blythe,

Coachella, and Calexico. This bioregion has four Indian reservations and several military bases. Imperial County is one of California's top agricultural counties. Cotton is a major crop.

Natural Systems

Salt marshes, freshwater ponds, and desert scrub are found in the Salton Sea National Wildlife Refuge.

- **Birds:** Great roadrunner, Gambel's quail, Albert's towhee, endangered Yuma clapper rail, egret, plover, northern pintail, Canada goose, snow goose, rough-legged hawk, peregrine falcon, tern, yellow-headed blackbird, hooded oriole, white-faced ibis.



Palm Canyon, Agua Caliente Cahuilla Indian Reservation

Restored wetlands in the Dos Palmas Preserve provide a desert oasis.

- **Birds:** Hooded oriole, warblers, snowy egret, osprey, American avocet.
- **Other animals:** Endangered desert pupfish, horned lizard.

Colorado River and surrounding land support many types of birds and wildlife.

- **Migratory birds:** Cormorant, merganser, white pelican, bald eagle.

Backcountry desert is home to many desert organisms.

- **Birds and mammals:** Desert bighorn sheep, feral burro, golden eagle, nesting prairie falcon, cactus wren, Gila woodpecker.
- **Endangered species:** Desert slender salamander, Yuma clapper rail, desert pupfish, razorback sucker (fish), unarmored threespine stickleback (fish), Coachella Valley fringe-toed lizard.

Factors Affecting Wildlife Diversity

- water management conflicts and water transfer
- inappropriate off-road vehicle use
- loss and degradation of dune habitats
- disruption of sand transport processes
- invasive plant and animal species
- growth and development of communities



Roadrunner with prey



Ocotillo and cholla cacti



Desert pupfish



Off-road driving

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Klamath/North Coast Bioregion

This bioregion features ancient redwood and Douglas fir forests. The rocky coastline has the state's wettest climate. Rainfall can be over 80 inches (203 cm)

per year. The coastal climate is cool, moist, and often foggy. Inland areas have hot, dry summers and low winter rainfall.

Human Social Systems


The largest cities are Redding and Eureka. Smaller cities include Clearlake, Ukiah, Arcata, Fort Bragg, Yreka, Mendocino, and Crescent City. The economy depends on cattle ranching, dairy farming, timber harvesting, and tourism. Fishing for Coho and king salmon is popular. The timber industry declined after unsustainable timber harvesting led to the listing of the northern spotted

owl as an endangered species, and the trend toward sustainable forest management grew.

Natural Systems

Wetlands along the coast provide food and shelter for many types of birds. Some live in California all year, and others stop to eat and rest during spring and fall migrations. Ancient redwood forests and other mature forests provide habitat to the northern spotted owl, an endangered species. The marbled murrelet is another endangered species found in these woods. A seabird, it nests in the tops of tall, old trees along the coast.

- **Mixed conifer forests:** White fir, Douglas fir, ponderosa pine, Sierra lodgepole pine, incense cedar, sugar pine, red pine, Jeffrey pine, mountain hemlock, knobcone pine, western redcedar, red alder, coast redwood, tanoak, Pacific madrone.



Del Norte Coast Redwoods State Park, California

- **Rare plants:** Burke's goldfield, Humboldt Bay owl's clover, coast lily, swamp harebell, Snow Mountain willowherb, marsh checkerbloom, pale yellow stonecrop, Scott Mountain phacelia, McDonald's rock cress, Klamath Mountain buckwheat, Oregon fireweed, Adobe lily.
- **Birds:** Bald eagle, California clapper rail, Aleutian Canada goose, osprey, Swainson's hawk, willow flycatcher, western sandpiper.
- **Other animals:** Deer, fox, black bear, mountain lion, Roosevelt elk, Pacific fisher (mammal), Point Arena mountain beaver, Oregon silverspot butterfly.
- **Rare animals:** Northern spotted owl, marbled murrelet, American peregrine falcon, Lotis blue butterfly, Trinity bristle snail, red-legged frog, Siskiyou Mountains salamander, Pacific fisher, Del Norte salamander, Karok Indian snail, wolverine, goshawk, Chinook salmon.
- **Endangered species:** Marbled murrelet, northern spotted owl, Humboldt Bay wallflower, western lily, Lost River sucker (fish), California clapper rail.



Old growth redwood forest, California



Roosevelt elk

Factors Affecting Wildlife Diversity

- water management conflicts
- in-stream gravel mining
- forest management can have either positive or negative influences. For example, thinning dense forests can enhance biodiversity.
- altered fire regimes
- habitat loss due to agriculture and community development
- livestock grazing
- invasive species

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Northern spotted owl



Logging



Modoc Bioregion

This area has the lowest human population of any of California's bioregions. The rural region features a variety of parks and wildlife refuges. Stretching across the Modoc Plateau, it

includes: forests, mountains, high desert, valleys, pine forests, and uplands created by volcanoes. Summers are hot and dry. Winters are cold and moist, with snow at higher elevations.

Human Social Systems

Modoc and Lassen counties are included in the Modoc Bioregion. The largest cities are Alturas, Susanville, Burney, and Magalia.

Two California Indian tribes—the Northern Paiute and the Paiute-Shoshone—live here. The economy depends on ranching and timber operations.

Natural Systems

The western side of the region features land created by volcanoes. Lassen Volcanic National Park is here, along with Lava Beds National Monument and several wildlife refuges. Mount Lassen last erupted in 1915. The eastern part of the bioregion is drier and features high desert ecosystems.

- **High desert vegetation:** Juniper and sagebrush communities.
- **Mountain forests:** Yellow pine, Jeffrey pine, white fir, mixed conifer, cedar, aspen.
- **Rare plants:** Yellow arrowleaf, balsam root, long-haired star tulip, spiny milkwort, Ash Creek ivesia, Raven's lomatium, woolly stenotus.
- **Birds:** Bald eagle, greater sandhill crane, osprey, Canada goose, black-crowned night heron, cinnamon teal, northern pintail, Swainson's hawk, sage grouse,



Mount Lassen, Lassen Volcanic National Park, California

hummingbird, great horned owl, goshawk, bank swallow.

- **Fish and other aquatic life:** Rainbow trout, Modoc sucker, Lost River sucker, Shasta crayfish.
- **Mammals:** Antelope, mule deer, pronghorn, marmot, black bear, coyote, porcupine, muskrat.
- **Endangered species:** Shasta crayfish, Modoc sucker, Lost River sucker, shortnose sucker.

Factors Affecting Wildlife Diversity

- livestock grazing
- feral horse grazing
- altered fire regimes
- western juniper expansion
- invasive plants
- forest management can have either positive or negative influences. For example, thinning dense forests can enhance biodiversity.
- water management conflicts and degradation of aquatic ecosystems

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Aspen and pine trees



Pronghorn antelope



Herding cattle



Mojave Bioregion

One of California's largest bioregions, the Mojave Bioregion's desert may look barren but actually is teeming with life. Streams and springs provide water that support many

forms of wildlife. This bioregion has the lowest elevation in North America, 282 feet (86 meters) below sea level in Death Valley National Park.

Human Social Systems

In the past, not many people lived in this desert region. Now the region supports rapidly growing cities. Palmdale is one such city. Others include: Victorville, Hesperia, Ridgecrest, and Barstow. The region also is home to three national parks, several

Indian reservations, and a variety of military operations. Mining for gold and other minerals is a major industry, along with ranching and livestock grazing.

Natural Systems

- **Habitats:** Desert wash, Mojave creosote bush, scattered desert saltbush, Joshua tree scrub, alkali scrub, palm oasis, juniper-piñon woodland, hardwood and conifer forests at higher elevations, cottonwood willow riparian forest, alkali marsh, open sandy dunes.
- **Rare plants:** White bear poppy, Barstow woolly sunflower, alkali mariposa lily, Red Rock poppy, Mojave monkeyflower, Stephen's beardtongue.



Mojave Desert, California

- **Birds:** Snowy plover, least sandpiper, killdeer, white pelican, teal, eagle, harrier, falcon, owls, migratory shorebirds, great blue heron, least Bell's vireo, red-tailed hawk, Canada goose.
- **Mammals:** Coyote, badger, kit fox, longtail pocket mouse, desert kangaroo rat, Merriam's kangaroo rat, black-tailed jackrabbit, bobcat.
- **Rare animals:** Mojave ground squirrel, prairie falcon, Le Conte's thrasher, Nelson's bighorn sheep, gray vireo (bird), desert tortoise, pale big-eared bat, Mojave tui chub (fish), cottontail marsh pupfish (found only in Death Valley National Park).
- **Endangered species:** California condor, Armargosa vole, Owens tui chub (fish), least Bell's vireo (bird).

Factors Affecting Wildlife Diversity

- multiple uses conflicting with wildlife on public lands
- habitat loss due to development
- groundwater overdrafting and loss of riparian habitat
- inappropriate off-road vehicle use
- excessive livestock, burro, and horse grazing
- invasive plants
- land management conflicts
- mining operations



Prickly pear cactus



Coyote



California condor



Gravel mining

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Sacramento Valley Bioregion

The Sacramento Valley Bioregion is a broad, flat valley. The Sacramento River drains from the Sierra Nevada Mountains and is the state's largest river. Most of the land is privately owned. Private landowners play an important conservation role in this bioregion. Over 75% of the known locations of 32 animal "species of concern" in California are on private land. (*Species of concern* is an informal designation for species for which there are some concerns regarding status and threats.)

Human Social Systems

More than 1.5 million people live in this region. Sacramento is the state capital. Other cities include: Redding, Chico, Davis, West Sacramento, and Roseville. Many people work for the state government. Others work in lobbying or public relations firms. Tomatoes, rice, olives, and other crops grow in Sutter, Yolo, and Colusa Counties. Food canneries, high-tech industries, and biotechnology contribute to the economy.

Natural Systems

The Sacramento Valley has hot summers, mild autumns, and foggy winters. Except during droughts, rainfall is frequent in winter. The weather usually is too warm for snow. The Sacramento



Sacramento National Wildlife Refuge, California

Valley is an important site for water birds in winter. More than 1.5 million ducks and 750,000 geese visit marshes along the Pacific Flyway in winter months.

- **Habitats:** Oak woodlands, riparian forests, vernal pools, freshwater marshes, grasslands.
- **Birds:** Northern pintail, snow goose, tundra swan, sandhill crane, mallard, grebe, peregrine falcon, heron, egret, hawk.
- **Other animals:** Black-tailed deer, coyote, river otter, muskrat, beaver, osprey, bald eagle, salmon, steelhead (fish), swallowtail butterfly.
- **Species of concern:** Swainson's hawk, burrowing owl, San Pablo vole, Buena Vista Lake shrew.
- **Endangered species:** Winter-run Chinook salmon, delta smelt (fish), giant garter snake, western yellow-billed cuckoo.

Factors Affecting Wildlife Diversity

- habitat loss due to urban, residential, and agricultural development
- water management conflicts and reduced water for wildlife
- water pollution
- invasive species

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Riparian habitat



Swainson's hawk



Housing development



Black-tailed deer



San Joaquin Valley Bioregion

Fertile soils and hot summers make this California's top agricultural region. As agriculture has grown, people have converted many of the region's

native grasslands, woodlands, and wetlands to farmland. Diversion of water for irrigation has dried up 95% of the streams. Public parks, reserves, and wildlife areas provide refuges for fish and wildlife. Summers are hot and dry with long, sunny days. Winters tend to be moist and foggy.

Human Social Systems

With more than 2 million people, the San Joaquin Valley ranks third among California's bioregions

in terms of human population. The largest cities are Fresno, Bakersfield, Modesto, and Stockton. Agriculture provides the basis for the economy. Oil and gas production also are important. The region includes several Indian reservations and one military operation.

Natural Systems

This region contains important oak woodlands and grasslands. The South Fork Wildlife Area supports 20% of California's remaining stream-bank cottonwood and willow forests. Natural grasslands, ponds, and marshes in the Tule Elk State Reserve support Tule elk. Four endangered species also live there. The Kern National Wildlife Refuge attracts peregrine falcons, ducks, shorebirds, and songbirds. Great blue herons, beavers, coyotes,



Oak trees, San Joaquin Valley, California

black bears, mountain lions, red-shouldered hawks, and mule deer make use of these habitats. These and other parks and wildlife refuges attract up to 1 million birds each winter, including bald eagles and a wide range of ducks.

- **Habitats:** Vernal pools, valley sink scrub, saltbush, freshwater marsh, grasslands, arid plains, orchards, and oak savannah.
- **Rare plants:** Hoover's woollystar, Mason's lilaepsis, San Joaquin woollythread, California hibiscus, valley elderberry.
- **Rare animals:** Western pond turtle, tricolored blackbird, northern harrier, western yellow-billed cuckoo, longhorn beetle.
- **Endangered species:** California tiger salamander, Swainson's hawk, giant kangaroo rat, Fresno kangaroo rat, San Joaquin kit fox, blunt-nosed leopard lizard, San Joaquin antelope squirrel, Tipton kangaroo rat.

Factors Affecting Wildlife Diversity


- habitat loss due to urban, residential, and agricultural development
- water management conflicts and reduced water for wildlife
- water pollution
- invasive species



Mountain lion and cub



Kit foxes



Grassland with poppies and lupines



Cattle in feed lot

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Sierra Bioregion

The Sierra Bioregion is named for the Sierra Nevada mountain range. Much of the state's water supply comes from these mountains. Mount Whitney is the country's highest peak (outside of Alaska). The region contains three national parks, eight national forests, and many other parks and recreation areas. These areas can host diverse forest types and provide habitat to a wide range of plant and animal species. Winters are snowy, and summers are dry and mild.

Human Social Systems

More than 650,000 people live in this region.

Rural towns and small cities include: Truckee, Placerville, Quincy, Auburn, South Lake Tahoe, and Bishop. Software and high-tech industries provide many jobs. Hydropower, tourism, recreation, logging, and cattle ranching also are important. Apple orchards and wineries are found in the foothills. Melting snow fills reservoirs. This provides about two-thirds of the water used in California for drinking, irrigation, and industry.

Natural Systems

Over half of California's plant species live in this region. The diverse habitats support two-thirds of the state's bird and mammal species. Half of the species of reptiles and amphibians in the state live in this region.



High meadow, Sierra Nevada, California

- **Habitats:** Alpine meadow, grassland, sagebrush, chaparral, forested riverbank.
- **Trees and shrubs:** Mixed conifer, blue oak, black oak, red fir, Jeffrey pine, ponderosa pine, bitter brush.
- **Birds:** Bald eagle, great grey owl, California spotted owl, northern goshawk, mountain chickadee, pine grosbeak, mountain quail, willow flycatcher.
- **Other animals:** California mountain king snake, California big horn sheep, mule deer, mountain lion, black bear, wolverine, mountain beaver, Pacific fisher, lodgepole chipmunk. Cold mountain streams are home to California Golden Trout, the state fish.
- **Endangered species:** Owens pupfish, Stebbins's morning-glory, Bakersfield cactus, blunt-nosed leopard lizard, California jewel-flower, California condor, Hartweg's golden sunburst (plant).



Bakersfield cactus, Bakersfield, California



Wolverine



Giant sequoias, California

Factors Affecting Wildlife Diversity

- dams, water diversions, and hydropower operations
- habitat loss due to community growth and land development
- altered fire patterns
- forest management can have either positive or negative influences. For example, it can thin dense forests and enhance biodiversity.
- livestock grazing
- invasive plants and introduced nonnative fish
- watershed fragmentation and fish barriers
- recreational pressures



Shasta Dam, California

Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



South Coast Bioregion

Landscapes in this bioregion range from flatlands to mountains. Ecosystems range from ocean to desert. The region contains two of California's largest cities: Los Angeles and San Diego. More

than in any other bioregion, urbanization has caused intense effects on natural resources. Urbanization in the South Coast Bioregion has resulted in loss of habitat, spread of nonnative species, and loss of native species.


Human Social Systems

More people live in this region than in any other part of California. The natural beauty and year-round mild climate make this a popular location. Recreation and tourism provide many jobs. Tourists visit Disneyland, Hollywood, Sea World,

and the San Diego Zoo. Tourism is important to Southern California communities. The miles of beaches and coastal mountain ranges are popular recreation spots. Many people work in the movie and television industries. Other important industries include: oil, shipping, banking, computers, and military. Agriculture and fishing also provide jobs.

Natural Systems

Many conservation projects are underway in this region. The Sespe Condor Sanctuary is home to the endangered California condor. Many coastal wetlands have been lost to development. However, some salt marshes and lagoons are being preserved or restored. Sage scrub communities also need protection. These communities support diverse kinds of wildlife. These include the California gnatcatcher, a threatened bird species.



Malibu Lagoon State Beach, Malibu, California

- **Lower elevation habitats:** Chaparral, juniper-pinyon woodland, grassland.
- **Mountain habitats:** Hardwood forest, southern oak, southern Jeffrey pine, southern yellow pine forest.
- **Rare plants:** San Diego barrel cactus, Conejo buckwheat, Plummer's mariposa lily, mountain springs bush lupine, Otay tarplant, Laguna Mountains jewelflower, San Jacinto prickly phlox, Mount Gleason Indian paintbrush.
- **Birds:** Hawks, heron, golden eagle, osprey, peregrine falcon, endangered brown pelican.
- **Other animals:** Mountain lion, coyote, black-tailed jackrabbit, grey fox, raccoon, mule deer.
- **Marine mammals:** Dolphins, whales, and California sea lion.
- **Rare animals:** Stephen's kangaroo rat, monarch butterfly, San Diego horned lizard, Peninsula desert bighorn sheep, orange-throated whiptail (lizard), California least tern, Belding's savannah sparrow, least Bell's vireo, Santa Ana sucker (fish), arroyo southwestern toad, Tehachapi pocket mouse.
- **Endangered species:** California least tern, Channel Islands foxes, California brown pelican, Santa Ana River woolly star, San Diego mesa mint, Nevin's barberry, bird-foot checkerbloom, California orcutt grass.

Factors Affecting Wildlife Diversity

- habitat loss due to urban growth and development
- water management conflicts and degradation of aquatic ecosystems
- invasive species
- altered fire regimes
- recreational pressures


Adapted from California Natural Resources Agency and California Department of Fish and Game, http://ceres.ca.gov/geo_area/bioregions/mapindex.html and <http://www.dfg.ca.gov/habitats/WDP/report.html>



Long-beaked dolphins



California brown pelican



La Verne hillside, California



Watering lawns